

Corn Silage Evaluation: MILK2000 Challenges & Opportunities With MILK2006

Professor Randy Shaver


Department of Dairy Science
University of Wisconsin - Madison
University of Wisconsin - Extension



Whole-Plant Corn Silage




Grain ~40-45% of WPDM

- Avg. 28% starch in WPDM
- Variable grain:stover



Stover= ~55-60% of WPDM

- Leaves = 15% of DM
- Stem = 20-25% of DM
- Cob+Shank+Husk= 20% of DM



80 to 98% starch digestibility

- Kernel maturity
- Kernel particle size
- Endosperm properties

40 to 70% NDFD

- lignin/NDF

NRC (2001) Dairy TDN


TDN 1-X =

$$***tdCP + (tdFA \times 2.25) + tdNDF + tdNFC - 7***$$

Why measure NDFD *in vitro* vs. calculating via lignin?

- Lignin wet chem assay difficult & its calibration with NIRS has been poor
- Lignin to NDFD equation is theoretically based
- Lignin explains only about half of the *in vitro* NDFD variation
 - Stover NDF & lignin contents ↑ & NDFD ↓ with maturity, while WP NDF & lignin contents are constant or ↓ as grain% increases

Relationship between lignified NDF and *in vitro* NDFD for corn & alfalfa forages



30-h NDFD (adapted from Allen, 2003) **vs. NDF digestibility**
calculated using NRC-01 lignin equation

<u>Whole-Plant Lignin, % DM</u>	<u>Calculated NDFD</u>	<u>30-h NDFD</u>
2.1	62	60
3.1	57	45
4.2	53	30

Measured NDFD or Estimated from Lignin?

NDF, %	Lignin, %	Calc. NDFD	30-h NDFD
45.0	3.52	56	46.0
45.0	3.26	57	48.4
45.0	3.32	57	54.4
45.1	3.18	57	55.0
45.0	3.43	56	67.3

- Corn silage data set from Van Amburgh (2004)
- Similar relationships from 36.5 to 51.8% NDF

Adapted from: Rick Grant, NRAES Silage Conf., 2006

The incubation time-point debate

- 48-hr.
 - Reflects maintenance intake for use in NRC summative equation
 - Less influenced by lag & rate, so possibly lower COV
- 30- or 24-hr.
 - 30-h more closely related to ruminal retention time
 - 30-h was used in most cow trials
 - Faster lab turn-around
 - Better lab efficiency at 24-h?


The incubation time-point debate

- MILK2000

- 48-h

- MILK2006

- 48-h default, with 30-h or 24-h User Defined Option
- Lab average NDFD required
 - NDFD DMIadjustment =
(avg. NDFD – NDFD) * 0.26
 - NDFD adjustment for summative TDN_{1x} equation



Adapted from Coors (data from Justen, 2004).

Variation in “normal “ corn silage NDF digestibility calculated using NRC-01 lignin equation and table data

<u>Whole-Plant Lignin, % DM</u>	<u>Calculated NDFD</u>
1.0 (2stdev)	65
1.8 (1stdev)	61
2.6 (avg.)	59
3.4 (1 stdev)	56
4.2 (2stdev)	56

NDFD -- MILK2000 vs. MILK2006

■ MILK2000


- A 1%-unit change in NDFD from lab average NDFD changes DMI 0.37 lb (Oba and Allen, 1999, JDS)
- Double counting of TDN & DMI changes related to changes in NDFD
 - Tine et al. (2001, JDS) and Oba and Allen (1999, JDS)
 - At production levels of intake, NDFD has minimal impact on NE_L content but does impact NE_L intake primarily thru its impact on DMI
- Calculation of NE_{L-3x} from TDN_{1x} as per NRC (1989)

■ MILK2006

- A 1%-unit change in NDFD from lab average NDFD changes DMI 0.26 lb (Jung, 2004, MN Nutr. Conf.; Oba and Allen, 2005, Tri-State Nutr. Conf.)
- NDFD used for calculating NE_{L-3x} adjusted for impact of NDFD on DMI (Oba and Allen, JDS, 1999)
- Calculation of NE_{L-3x} from TDN_{1x} via DE and ME as per NRC (2001)


Corn Silage NDFD% vs. NE_{L3x}

Calculated assuming corn silage with 35% DM proc., 27% starch, 45% NDF, and 58% avg. NDFD



Corn Silage NDFD% vs. Milk per Ton

Calculated assuming corn silage with 35% DM proc., 27% starch, 45% NDF, and 58% avg. NDFD



NRC (2001) Dairy TDN

TDN 1-X =

tdCP + (tdFA x 2.25) + tdNDF + **tdNFC** -7

NRC (2001) Dairy TDN


$$tdNFC = NFC\% \times 0.98 \times PAF$$

	<u>PAF</u>
Corn grain, ground dry	1.00
Corn grain, ground high moisture	1.04
Corn silage, normal	0.94
Corn silage, mature	0.87


Schwab-Shaver Energy Equation

$\text{TDN}_{1-x} =$

$\text{DIG}_{\text{CP}} + \text{DIG}_{\text{FA}} + \text{DIG}_{\text{Starch}} + \text{DIG}_{\text{NSTNFC}} + \text{DIG}_{\text{NDF}} - 7$



Predicted Starch Digestibility




Adapted from Schwab et al., 2003.

Differences in calculation of *tdNFC*

Based on corn silage with 41% NFC & 28% starch


<u>WP DM %</u>	<u>NRC-01 <i>tdNFC</i>%</u>	<u>Schwab et al., 2003 <i>tdStarch&NSTNFC</i>%</u>	
		Unprocessed	Processed
30	40	40	40
35	38	39	40
40	35	36	39
45	35	34	38

Corn Silage WP DM% vs. TDN_{1x}



Corn Silage WP DM% vs. Milk per Ton

Calculated assuming corn silage with 27% starch, 45% NDF, and 58% NDFD



Evaluating Starch Digestion in Ruminants.....

**In Vivo..... Total Tract Collections & Digesta Markers
Cannulae (Rumen, Duodenum, Ileum)**

**Macro In Situ.... Rumen cannulae
Incubation time?
Starch-Feeds cannot be fine ground
Post-Ruminal Enzymatic?**

**In Vitro.... Rumen fluid
Incubation time ?
Starch-Feeds cannot be fine ground
Post-Ruminal Enzymatic?**

Corn Silage Processing Score

Mertens, USDFRC & Dairyland Labs, Arcadia, WI

- Ro-Tap Shaker
 - 9 sieves (0.6 thru 19 mm) and pan
 - Analyze for starch on 4.75 mm & greater sieves

% of starch passing

4.75 mm sieve

>70%

70% to 50%

< 50%

CSPS

Optimum

Average

Poor

Kernels and Large Fragments Were Retained on > 4.75 -mm Sieves




USDA-ARS

US Dairy Forage Research Center

Corn Silage KPS vs. TDN_{1x}

Calculated assuming corn silage with 40% DM 27% starch, 45% NDF, and 58% NDFD





MILK2006

Degree of Starch Access (DSA)

Blasel, Hoffman and Shaver, JAFST, 2006


- Adaptation of food industry assay “Degree of Starch Gelatinization”
- Detects particle size, moisture, and vitreousness differences in corn samples
- Appears to offer better characterization of processed corn silage samples than KPS
- DSA can be related to total tract starch digestion
 - More animal validation data needed
- Pilot study of assay across labs in progress






Blazel et al., 2005

DSA vs. Total Tract Starch Digestibility from Literature Sources



Corn Silage DSA vs. TDN_{1x}

Calculated assuming corn silage with 27% starch, 45% NDF, & 58% NDFD



MILK2006: Starch Digestion


User Defined Options

- Default
 - WP DM & Kernel Processing Regressions
- KPS
- DSA
- Ruminant in situ plus post-ruminant in vitro

TDN_{1x} Simulation -- Input Extremes

LQ 45% DM unproc., 20% starch, 54% NDF, and 46% NDFD


HQ 30% DM proc., 34% starch, 36% NDF, and 70% NDFD



Milk per Ton Simulation -- Input Extremes

LQ 45% DM unproc., 20% starch, 54% NDF, and 46% NDFD


HQ 30% DM proc., 34% starch, 36% NDF, and 70% NDFD



Milk per Ton -- High NDF, NDFD vs. Low NDF, NDFD Ivan et al., JDS, 2005


LF 26% starch, 49% NDF, 58% NDFD

HF 22% starch, 53% NDF, 67% NDFD




Milk per Ton -- High NDF, NDFD vs. Low NDF, NDFD

Ivan et al., JDS, 2005




Treatment differences for model-predicted milk per ton versus milk per ton from in vivo data



Calculated from 10 JDS papers with 13 comparisons

Treatment differences for model-predicted milk per day versus milk per day in vivo data



Calculated from 10 JDS papers with 13 comparisons

UW Correlations

n = 3727 treatment means


r-values	Milk per Ton DM 2006	Milk per Ton DM 2000	Milk per Ton DM 1995	Milk per Ton DM 1991
NDF	-0.46	-0.40	-0.94	-0.99
Starch	0.48	0.44	0.75	0.74
NDFD	0.49	0.70	0.16	-0.10
StarchD	0.30	0.21	-0.25	-0.27

UW Correlations

n = 3727 treatment means

r-values	Milk per Acre 2006	Milk per Acre 2000	Milk per Acre 1995	Milk per Acre 1991
DM Yield tons/acre	0.97	0.97	0.88	0.85
Milk per Ton DM	0.23	0.20	0.52	0.61

Relationship between milk per acre and milk per ton of corn hybrids in South Central WI during 2002.



Visit UW-Madison Dairy Science Department's Website

<http://www.wisc.edu/dysci/>



**EXCELLENCE IN
EDUCATION AND DISCOVERY**

UNIVERSITY OF WISCONSIN - MADISON

www.wisc.edu/dysci



THE UNIVERSITY
of
WISCONSIN
MADISON